

The Relationship between Saccadic Suppression and Perceptual Stability

Tamara L. Watson^{1,*} and Bart Krekelberg¹

¹Center for Molecular and Behavioral Neuroscience
Rutgers University
197 University Avenue
Newark, NJ 07102
USA

Summary

Introspection makes it clear that we do not see the visual motion generated by our saccadic eye movements. We refer to the lack of awareness of the motion across the retina that is generated by a saccade as saccadic omission [1]: the visual stimulus generated by the saccade is omitted from our subjective awareness. In the laboratory, saccadic omission is often studied by investigating saccadic suppression, the reduction in visual sensitivity before and during a saccade (see Ross et al. [2] and Wurtz [3] for reviews). We investigated whether perceptual stability requires that a mechanism like saccadic suppression removes perisaccadic stimuli from visual processing to prevent their presumed harmful effect on perceptual stability [4, 5]. Our results show that a stimulus that undergoes saccadic omission can nevertheless generate a shape contrast illusion. This illusion can be generated when the inducer and test stimulus are separated in space and is therefore thought to be generated at a later stage of visual processing [6]. This shows that perceptual stability is attained without removing stimuli from processing and suggests a conceptually new view of perceptual stability in which perisaccadic stimuli are processed by the early visual system, but these signals are prevented from reaching awareness at a later stage of processing.

Results

To determine the fate of a saccadically omitted stimulus, we used a visual shape illusion in which the presentation of a line distorts the perceived shape of a subsequently presented ellipse [6] (Figure 1). Our goal was to present the line within 75 ms preceding a saccade—such that the observer would not be aware of it on some trials—and then determine whether such a saccadically omitted stimulus nevertheless retained the ability to change the subsequent perceived shape of an ellipse. The observers first reported whether or not they saw the line, which could be horizontal or vertical and was physically present on only 50% of trials. They then reported the shape of the ellipse that was presented once the eyes had landed. This allowed us to determine the percentage of trials in which they perceived the ellipse to be horizontally elongated. We refer to this as %PHE (perceived horizontal elongation). To quantify the size of the illusion, we presented ellipse stimuli with a physical horizontal or vertical elongation of 10%, 7.5%, 5%, or 2.5% of the diameter (3° of visual angle). A separate cumulative

Gaussian was fitted to the %PHE in the horizontal and vertical line conditions. The subjective point of circularity was taken as the physical elongation at which the fitted function reached 50%PHE and the strength of the illusion was defined as the difference between the points of subjective circularity induced by presentation of the horizontal and vertical lines. Subjective circularity is measured in units of percentage of elongation of the diameter of the test circle. We first confirmed that the shape illusion occurs at fixation for briefly flashed stimuli (Figure 2 and Supplemental Results, Experiment 1, available online) and also when an eye movement intervenes in the 250 ms between the presentation of the line and the ellipse (Supplemental Results, Experiment 2). We then analyzed separately the saccade trials in which the subject reported no subjective awareness of the inducing line (saccadic omission was achieved) and those trials in which subjects reported awareness of the inducing line (saccadic omission was not achieved).

Figure 2 shows our critical finding that the participants had a significant shape illusion even without awareness of the inducer ($t_4 = 2.8125$, $p < 0.05$ [see Supplemental Results for further details]). Hence, even when saccadic omission was complete, the omitted stimulus affected subsequent perception. This implies that the line was processed at least by those higher visual areas responsible for this shape illusion (see Discussion) and shows that perceptual stability does not require perisaccadic visual stimuli to be removed from processing.

The data in Figure 2 also show that the shape illusion was reduced for those lines that were reported not to be seen compared to lines that were reported to be seen. This shows that the influence a stimulus exerts on visual processing is diminished most when it is not seen. Our data show, however, that this suppression is not enough to explain the omission because a stimulus that is only partially suppressed nevertheless can be fully omitted.

Our interpretation of the data relies on taking seriously the subjective report and its binary nature (seen/not seen). We believe that this is appropriate and in fact essential in this context because under most everyday circumstances, normal observers do not perceive any retinal motion during an eye movement and the absolute nature of this phenomenon requires an explanation. This notwithstanding, there could be a concern that, in the critical condition of complete omission with incomplete suppression, observers may have experienced the stimulus despite answering to the contrary. In the Supplemental Data we analyzed this possibility and conclude that to generate an effect of the appropriate size the observers would have to be wrong about their perceptual experience in more than 25% of trials. We believe this to be an unlikely explanation of our findings. Moreover, questioning upon completion of the experiments also reassured us that participants were reporting not seeing the stimulus when they were convinced that no stimulus existed, i.e., when they were experiencing saccadic omission.

Discussion

Our data show that a stimulus that is successfully removed from awareness by saccadic omission is nevertheless processed by

*Correspondence: tamara@vision.rutgers.edu




Figure 1. The Shape Contrast Illusion
A circle presented after a line appears elongated orthogonal to the line orientation.

the visual system. This shows that perceptual stability can be attained without removing a perisaccadic stimulus from visual processing. We discuss this finding in light of a number of debates surrounding perceptual stability and saccadic suppression.

Early versus Late

The shape contrast illusion we used here has been shown to survive when the inducing and test stimuli are separated in space, indicating that the illusion involves an area whose shape-selective neurons have large receptive fields such as superior temporal sulcus or inferotemporal cortex [6]. Our data therefore suggest that saccadically omitted stimuli are nevertheless processed by such higher visual areas. The fact that the illusion is reduced in size during omission suggests that there may be a (possibly early) stage at which the efficacy of a stimulus is reduced [4]. A modulation, including suppression, of the visual responses of single cells has been shown as early as the lateral geniculate nucleus [7].

Retinal versus Extraretinal

Two main mechanisms of saccadic suppression have been proposed. The first relies on an extraretinal signal that prepares the visual system to discount the upcoming high velocity perisaccadic retinal motion. The second is a form of masking whereby the perisaccadic stimulus is “wiped out” by the stable, fixated stimuli before or after the saccade. Both of these mechanisms are thought to occur early in the visual processing hierarchy to stabilize the visual world by removing the intrusion of the perisaccadic stimulus into the stable representation of fixated scenes [3]. The extraretinal component of saccadic suppression suppresses mainly low-spatial-frequency, luminance-defined stimuli [4], which are considered the domain of the magnocellular visual pathway, and does not appear to affect judgments involving isoluminant chromatically defined stimuli, which are processed mainly by the parvocellular pathway [4].

In our experiments the inducing line was presented just before the eye movement, a time at which the extraretinal mechanism of suppression is known to operate. Moreover, the luminance-modulated line was flashed briefly and would likely activate the magnocellular pathway. Masking mechanisms will also be activated by the background luminance of




Figure 2. Average Strength of the Shape Contrast Illusion

Illusion strength is measured as the difference in point of subjective circularity of test stimuli following presentation of a horizontal compared to a vertical line (see main text). This calculation was carried out separately during fixation (Baseline), when the inducer was seen during an eye movement (Inducer Seen), and when it was not seen during an eye movement (Inducer Not Seen). Error bars represent 95% confidence intervals.

the CRT screen and by presentation of the ellipse after the eye movement. Hence, our experiments do not allow us to distinguish between extraretinal and masking mechanisms of suppression, but it is likely that both mechanisms were operating and therefore both were only partially effective in removing the stimulus from processing even when omission was complete.

Dorsal versus Ventral

The shape contrast illusion reported here presumably arises from interactions in the ventral stream. Hence, one could hypothesize that complete omission with incomplete suppression is a phenomenon that is unique to the ventral stream. Or, stated differently, that the dorsal stream undergoes suppression and omission, while the ventral stream only undergoes omission. Such a distinction is consistent with behavioral data that show that saccadic suppression mainly targets stimuli that drive the magnocellular stream, which, in turn, mainly projects to the dorsal stream. The neural evidence, however, suggests that saccadic reductions in firing rate can be found in both magnocellular and parvocellular cells of the lateral geniculate nucleus [7] and in both dorsal [8, 9] and ventral [10] cortical areas. Hence, it seems unlikely that ventral areas undergo no suppression. Conversely, it is also unlikely that suppression is complete in dorsal areas; the perisaccadic reversal of direction preference reported by Thiele et al. [8], for instance, suggests that saccadic suppression is more complex than simply not responding to perisaccadic stimuli.

Our data, together with the extensive literature on perisaccadic changes in neural response properties, lead us to rethink the purpose of saccadic suppression. We speculate that perisaccadic signals are useful; the visual motion signals generated by the eye movement, for instance, are excellent indicators of the size and speed of the eye movement. These signals could be used to improve perceptual stability as long as they do not directly enter visual awareness. This suggests that perisaccadic processing leading to saccadic omission consists of three conceptually different components: a (possibly early visual) reduction in sensitivity (saccadic suppression) and a component that processes perisaccadic signals to extract information useful for visual stability (saccadic information extraction). Additionally, perceptual stability requires a conceptually separate

component that prevents the perisaccadic signals from reaching awareness (saccadic omission).

Most research whose purported goal is to investigate the neural mechanisms of saccadic omission has only looked for signatures of the first component. Even though some studies did indeed find neural correlates of a reduced perisaccadic sensitivity [11–15], others have reported complex changes in perisaccadic activity that are difficult to reconcile with a mere suppression of visual processing [7–9, 15–17] (see [3] for a review). Within our framework, however, complex changes in activity are expected because the visual system tries to extract information from the perisaccadic visual inputs or because the system processes the information while keeping it hidden from awareness. Importantly, our data show that even if a neural correlate of the change in sensitivity could be found in early visual areas, then we would still not understand the neural basis of saccadic omission and how it leads to perceptual stability. Our conceptualization, therefore, provides a useful framework to guide further research and interpret findings about the neural basis of perceptual stability.

Experimental Procedures

Subjects

All conditions were completed by four naive participants and one experimenter. The four naive participants received remuneration and all had normal or corrected to normal vision.

Visual Stimuli

The stimuli consisted of a bar and a ring. Both were presented with half cosine profile luminance graduated edges against a background luminance of 45 cd/m² and a peak luminance of 47 cd/m². The inducing line was 7° × 1° of visual angle, whereas the circle had a diameter of 3° of visual angle and an outline line width of 0.8° of visual angle. The bar was present on the screen for 16 ms, and the ring was present for 100 ms. The ring was always presented 250 ms after the onset of the inducing line.

Apparatus

Stimuli were presented on a Sony FD Trinitron (GDM-C520) CRT monitor with a resolution of 1024 × 768 pixels at a refresh rate of 120 Hz. Stimuli were generated with Neurostim (<http://neurostim.sourceforge.net>). Eye movements were measured with a head-mounted Eyelink II eye tracker (SR Research, Mississauga, Canada). The pupil of the left eye was tracked at a sample rate of 500 Hz and a spatial resolution of 0.1°. Participants were seated in a darkened room at a 57 cm distance from the display. Head movements were restricted via individually molded bite bars.

Procedure

Experiment 1

For testing the basic shape contrast illusion with stimuli optimized for saccadic suppression, experiment 1 was carried out without eye movements and all stimuli were presented at the center of the screen. The inducing bar could be either horizontal or vertical but was only presented on half of all trials while the test stimuli were presented with a horizontal or vertical elongation of 10%, 7.5%, 5%, and 2.5% of the diameter. All possible pairings of bar and ring orientation were presented in a randomized order and 40 trials were collected for each, making a total of 640 trials.

Participants were made aware that only half the trials contained a line stimulus and were asked to report the orientation of the elongation of the ring stimulus via a key press once the stimulus had appeared.

Experiment 2

Stimuli were as described in experiment 1 with the exception of the bar being presented 5° above the midpoint of the monitor and the ellipse being presented 5° below the midpoint of the monitor.

In this experiment, if at any stage fixation was not achieved or maintained as required, the trial was terminated. The terminated trial was repeated at a later stage within the same test session. Participants were first shown a 0.2° × 0.2° white fixation point 5° above the center of the screen. They were required to fixate on this until it disappeared. One thousand milliseconds after fixation was achieved, a 0.3° × 0.3° white saccade target appeared 5° below the center of the screen. The cue to move the eyes (disappearance

of the current fixation point) was given between 1900 to 2200 ms after initial fixation. At 2200 ms after initial fixation was achieved the inducing bar appeared around the location of the now invisible upper fixation point. This timing aimed to present the inducing bar to the participant within the window of saccadic suppression but before the eye had started the saccade. The ring stimulus was then presented at the location of the saccade target 250 ms after presentation of the inducing bar. Participants were required to have achieved fixation of this target at least 100 ms prior to presentation of the ring stimulus. This fixation requirement ensured that the ring stimulus always landed on a stationary retina.

Participants reported whether they saw the inducing bar and the axis of elongation of the ring. Both tasks were carried out via a key press after the completion of the eye movement.

An additional set of trials were carried out with the same procedure; however, the inducing line was presented 2050 ms after initial fixation was achieved. This ensured that the inducing line was presented well before the saccadic suppression window and provided an estimate of participants' miss rate for the inducing line.

Data Analysis

Experiments 1 and 2

A separate cumulative Gaussian distribution was fit to the reported perceived axis of elongation of the ellipse during trials with a vertical inducer, trials with a horizontal inducer, and trials without an inducer. The Psignifit toolbox for Matlab [18, 19] was used for this analysis. The threshold at 50% was considered the point of subjective circularity under each different inducing condition. As a test of significance, we used the bootstrapping methods of the Psignifit toolbox to test the null hypothesis that the cumulative Gaussians for vertical and horizontal bars could be generated by the same underlying distribution.

Experiment 2

Trials during which appropriate fixation was not achieved were automatically discarded. Additionally, we selected only those trials in which the test stimulus was presented during the 75 ms before an eye movement was initiated. Additional to the separation of trials according to orientation of the inducing bar, trials were also grouped according to participants' responses about its visibility.

Supplemental Data

Supplemental Data include Supplemental Results and five figures and can be found with this article online at [http://www.cell.com/current-biology/supplemental/S0960-9822\(09\)01051-3](http://www.cell.com/current-biology/supplemental/S0960-9822(09)01051-3).

Acknowledgments

The authors are grateful to Kai Schreiber and Adam Morris for comments on the manuscript and thank the Human Frontiers Science Program (T.L.W.) and the Pew Charitable Trusts (B.K.) for financial support.

Received: March 10, 2009

Revised: April 10, 2009

Accepted: April 20, 2009

Published online: May 28, 2009

References

1. Campbell, F.W., and Wurtz, R.H. (1978). Saccadic omission: Why we do not see a grey-out during a saccadic eye movement. *Vision Res.* 18, 1297–1303.
2. Ross, J., Morrone, M.C., Goldberg, M.E., and Burr, D.C. (2001). Changes in visual perception at the time of saccades. *Trends Neurosci.* 24, 113–121.
3. Wurtz, R.H. (2008). Neuronal mechanisms of visual stability. *Vision Res.* 48, 2070–2089.
4. Burr, D.C., Morrone, M.C., and Ross, J. (1994). Selective suppression of the magnocellular visual pathway during saccadic eye movements. *Nature* 371, 511–513.
5. Holt, E.B. (1903). Eye movements and central anaesthesia. *Psychol. Rev.* 4, 3–45.
6. Suzuki, S., and Cavanagh, P. (1998). A shape-contrast effect for briefly presented stimuli. *J. Exp. Psychol. Hum. Percept. Perform* 24, 1315–1341.

7. Reppas, J.B., Usrey, W.M., and Reid, R.C. (2002). Saccadic eye movements modulate visual responses in the lateral geniculate nucleus. *Neuron* 35, 961–974.
8. Thiele, A., Henning, P., Kubischik, M., and Hoffmann, K.P. (2002). Neural mechanisms of saccadic suppression. *Science* 295, 2460–2462.
9. Ibbotson, M.R., Crowder, N.A., Cloherty, S.L., Price, N.S., and Mustari, M.J. (2008). Saccadic modulation of neural responses: Possible roles in saccadic suppression, enhancement, and time compression. *J. Neurosci.* 28, 10952–10960.
10. Tolias, A.S., Moore, T., Smirnakis, S.M., Tehovnik, E.J., Siapas, A.G., and Schiller, P.H. (2001). Eye movements modulate visual receptive fields of V4 neurons. *Neuron* 29, 757–767.
11. Bartlett, J.R., Doty, R.W., Lee, B.B., Sr., and Sakakura, H. (1976). Influence of saccadic eye movements on geniculostriate excitability in normal monkeys. *Exp. Brain Res.* 25, 487–509.
12. Duffy, F.H., and Burchfiel, J.L. (1975). Eye movement-related inhibition of primate visual neurons. *Brain Res.* 89, 121–132.
13. Sylvester, R., Haynes, J.D., and Rees, G. (2005). Saccades differentially modulate human LGN and V1 responses in the presence and absence of visual stimulation. *Curr. Biol.* 15, 37–41.
14. Vallines, I., and Greenlee, M.W. (2006). Saccadic suppression of retinotopically localized blood oxygen level-dependent responses in human primary visual area V1. *J. Neurosci.* 26, 5965–5969.
15. Price, N.S., Ibbotson, M.R., Ono, S., and Mustari, M.J. (2005). Rapid processing of retinal slip during saccades in macaque area MT. *J. Neurophysiol.* 94, 235–246.
16. Kleiser, R., Seitz, R.J., and Krekelberg, B. (2004). Neural correlates of saccadic suppression in humans. *Curr. Biol.* 14, 386–390.
17. Sylvester, R., and Rees, G. (2006). Extraretinal saccadic signals in human LGN and early retinotopic cortex. *Neuroimage* 30, 214–219.
18. Wichmann, F.A., and Hill, N.J. (2001). The psychometric function: II. Bootstrap-based confidence intervals and sampling. *Percept. Psychophys.* 63, 1314–1329.
19. Wichmann, F.A., and Hill, N.J. (2001). The psychometric function: I. Fitting, sampling, and goodness of fit. *Percept. Psychophys.* 63, 1293–1313.

Supplemental Data

The Relationship between Saccadic

Suppression and Perceptual Stability

Tamara Watson and Bart Krekelberg

Supplemental Results

Experiment 1

As a test of significance, we used the bootstrapping methods of the Psignifit toolbox [1,2] to test the null hypothesis that the cumulative Gaussian fitted to the %PHE (Perceived Horizontal Elongation) in the horizontal and vertical line conditions could be generated by the same underlying distribution. All participants showed a significant difference in point of subjective circularity depending on the orientation of the inducing bar both before and after completion of experiment two ($p < 0.001$ for each participant). Four participants displayed the classic shape contrast effect while one participant showed a consistently opposite, attractive effect (Figure S1). As we were interested in whether the illusion remains despite saccadic omission we only measured the relative difference in illusion between the fixation and eye movement conditions and the sign of the illusion here does not concern us.

All participants except the individual who had an attractive illusion showed a reduction in the size of the illusion after the substantial practice gained during the intervening eye movement trials.




Figure S1. Strength of the shape contrast illusion. Individual participants illusion strength at baseline (dark bars), when the inducer is seen during an eye movement (mid bars)

and not seen during an eye movement (light bars). Error bars represent 95% confidence intervals.

Experiment 2

Omission

To estimate the effectiveness of saccadic omission, we compared the number of inducers reported as unseen in two time windows. The first contained the bars presented between 200 to 80 ms before the saccade; this is expected to be outside the saccadic omission window [3]. The second window contained the bars presented within the omission window (75 ms before until saccade onset). On average each participant reported less than 10% of the inducers as unseen when they were presented in the first window while almost 50% were reported as unseen in the omission window (Figure S2).




Figure S2. Stimulus miss rate during eye movements. Individual participants miss rate for inducer presented before the saccadic suppression window (dark bars) and within the suppression window (light bars). All subjects show clear evidence of saccadic omission in the relevant time window.

Illusion

The shape illusion in the unseen condition was individually significant in three out of five subjects. An example of the cumulative Gaussian fits to one naïve participant's data in each condition of experiment two can be seen in Figure S3, demonstrating the size of the illusion when no inducing line was presented and when the inducer was omitted or visible. Figure S4 shows, for the same participant, the time of onset of the inducing line relative to saccade onset when the line stimulus was reported as omitted or visible. It is notable that omission occurs most readily when the stimulus was presented as close as possible to saccade onset. This is consistent with previous studies showing that peri-saccadic suppressive effects increase from 75ms before and are maximal at saccade onset [3]. Individually, participant two rarely reported the line stimulus as 'not seen' (Figure S2). As a consequence it was exceedingly difficult to gather enough trials in the 'not-seen' condition to estimate the strength of the illusion. This is reflected in this subject's large confidence intervals (figure S1). Subject 5 had an illusion

that was small at fixation and non-significant during eye movements, irrespective of whether the line was seen or unseen, hence this subject's results have no direct bearing on the hypothesis we tested. These individual differences notwithstanding, we found that at the group level the effect was significant. We performed a t-test comparing the two points of subjective equality depending on the orientation of the inducer. The inducer orientation had a significant effect both for seen ($t_4=4.2519$, $P<0.05$) and saccadically omitted ($t_4=2.8125$, $P<0.05$) oriented lines.




Figure S3. Gaussian fits to Experiment Two data for one naive participant; Blue = Vertical Inducer, Green = horizontal inducer. Test stimulus shape is represented in units of percentage of elongation of the 3 degree of visual angle diameter of a test circle (the circle is represented by 0 on the X axis). A shows the data when the inducing lines were assigned an orientation but their contrast was set to zero (i.e they were not displayed- this graph serves as a visual indicator of the typical noise in our behavioral measurements). B; the inducer was present but reported as unseen. C; the inducer was present and reported as seen.




Figure S4. Time of onset of the inducing line relative to saccade onset at time 0. Data is from naïve participant shown in figure S3 above. A; Inducer reported as unseen. B; Inducer reported as seen.

Given that it is possible that subjects erroneously reported lines as ‘unseen’, we analyzed how many such mistakes an observer would have to make to generate an illusion of the observed size. Data from experiment 2 were re-sampled to simulate the effect of responding ‘unseen’ in a trial that was actually seen. Trials were chosen at random from those correctly identified as containing an inducer to replace trials where the inducer was correctly identified as not presented. The percentage of trials replaced in this way was determined by each participant’s own miss rate from trials where the inducer appeared well before the eye movement and was therefore expected to be visible (Figure S2). As an even more stringent test we also simulated an insertion rate of 25% for all participants as a comparison. This re-sampling procedure was repeated 1000 times at each insertion rate and the mean size of the illusion was recorded. The data shown in Figure S5 suggest that even when trials containing correctly detected inducers are misreported as ‘unseen’ much more than 25% of all trials would need to be under reported in this way to generate an illusion of the size found during saccadic omission. At a 25% insertion rate only one participant matched the strength of illusion found during saccadic omission.




Figure S5. Strength of the illusion assuming mistaken reporting. Individual participants observed illusion strength during omission (dark bars), estimated illusion strength at the measured error rate during an eye movement (mid bars) and estimated illusion strength at a 25% error rate (light bars). Error bars represent 95% confidence intervals.

Supplemental References

1. Wichmann, F.A., and Hill, N.J. (2001). The psychometric function: II. Bootstrap-based confidence intervals and sampling. *Percept Psychophys* 8, 1314-29.
2. Wichmann, F.A., and Hill, N.J. (2001). The psychometric function: I. Fitting, sampling, and goodness of fit. *Percept Psychophys* 8, 1293-313.
3. Diamond, M.R., Ross, J., and Morrone, M.C. (2000). Extraretinal control of saccadic suppression. *J Neurosci* 9, 3449-55.